

The Positive Perspectives Survey Report

**A view into the lives
of people living with HIV**





Contents

In order to allow you to explore this report as easily as possible, we have implemented a simple navigation.

This contents page will take you to the relevant part of the report and the ViiV Healthcare logo will return you to this point.

	Page
Glossary	3
Foreword	4
About this report	5
Key insights	6
Expert panel	8
Survey methodology	9
Diagnosis	10
Stigma and disclosure	16
Getting on treatment	22
Treatment satisfaction	23
Conversations between PLHIV and their doctor	25
Long-term effects of treatment	26
References	28
About ViiV Healthcare	29

There are a number of terms or abbreviations (acronyms) that may be unfamiliar. They are listed in the table below.

Antiretroviral Therapy (ART)	Medications used to treat or prevent HIV; can reduce the amount of virus in blood to undetectable levels, preventing HIV-related illness or transmission.
Bisexual	Sexually attracted to both men and women
Comorbidity	A condition that exists at the same time as another condition
Heterosexual	Sexually attracted to people of the opposite sex
Homosexual	Sexually attracted to one's own sex
MSM	Men who have sex with men
PLHIV	People living with HIV
QoL	Quality of life
Switch	To change from one treatment to another
WSW	Women who have sex with women

Survey respondent numbers:

Throughout this report you will see that alongside every percentage/statistic from the Positive Perspectives survey, there is a number in brackets. This is the actual number of respondents represented by that percentage. For example, if a figure is shown as 68% – which reflects 755 out of 1,111 PLHIV – then it would be shown as: 68% (755).

Please refer to page 9 for the base values of country respondents and demographic breakdowns.

Glossary

Foreword

There have been incredible advances in biomedical interventions for the treatment and prevention of HIV, and current treatments are highly effective in preventing disease progression, death from AIDS, and transmission to others. However, in order to provide a truly holistic package of care for PLHIV, it is vital that these clinical advances are complemented with a strong understanding of the perspectives of PLHIV, including an understanding of the pressures that stigma and disclosure decisions put on PLHIV at diagnosis, and throughout their lives.

The importance of involving experiences and knowledge of PLHIV in decision-making around their medical care has been appreciated ever since it was enshrined in the 1983 Denver Principles. With this background, the Positive Perspectives survey sought to capture the experiences and opinions of a large and diverse international group of PLHIV, on the topics of the psychosocial aspects of living with the virus, dialogues between patient and healthcare provider and satisfaction with current antiretroviral therapies.

Regarding their relationship with HCPs, nearly all (94%) of those surveyed said that they felt quite or very comfortable in raising concerns with their HCPs. For me, this captures the essence of why we conducted this research. The willingness among PLHIV to discuss these issues with caregivers tells us loud and clear that we need to be ready to support this community, not only with medical treatment, but with counsel on every aspect of living with HIV, and activities which help them where they feel the greatest need.



By Dr Benjamin Young

Senior Vice President and Chief Medical
Officer of the International Association
of Providers in AIDS Care (IAPAC)



About this report

This report focuses on the results from the Positive Perspectives survey that was created by ViiV Healthcare to understand the emerging needs of PLHIV.

This report covers:

- The emotional support/guidance PLHIV received at diagnosis and where they turn to for support now
- How open PLHIV are about disclosing their status
- Experiences with stigma
- PLHIV and their treatment
- Enhancing communications between PLHIV and their doctor



Key insights

25%

(283) of those who were surveyed felt that **better doctor education** will reduce feelings of stigma^{1,2}



Stigma and disclosure

82%

(909) of PLHIV surveyed have experienced a **form of stigma related to their HIV in the last 12 months**^{1,2}



Getting on treatment

52%

(571) of those surveyed were **taking ART within six months** of their HIV diagnosis³



Diagnosis

68%

(759) of those surveyed indicated that they were **offered support**, referred for counselling or informed about support services by their HIV doctor **at diagnosis**^{1,2}



43%

(355) of those that have **switched did so in the last year** with side effects being the predominant reason^{3,4}

Key insights

Conversation between PLHIV and their doctor



89%

(987) of PLHIV believe that advances in HIV treatment **will improve their quality of life**^{1,2}

Treatment satisfaction



72%

(783) worry about the **long-term effects of their HIV treatment**³



56%

(604) of those surveyed that are **currently on treatment** are **very satisfied with it**³

Expert panel

The Positive Perspectives survey was conducted by ViiV Healthcare in collaboration with an international, multi-disciplinary expert panel that included: HIV physicians, PLHIV and patient group representatives. The expert panel contributed to the development of the survey themes and quantitative research questions, and participated in the communication of these results.



Brent Allan
Australia

Senior Advisor - Australasian Society for HIV Medicine (AUS) & International Council of AIDS Service Organisations (CDN)



Diego Garcia Morcillo
Spain

Member of the European AIDS Treatment Group (EATG), European Community Advisory Board (ECAB) and Chair of the BOD of ADHARA



Professor Rob Horne
UK

Professor of Behavioural Medicine at the School of Pharmacy, University College London and Founder of Spoonful of Sugar Ltd



Moritz Krehl
Germany

Member of the European AIDS Treatment Group; and scholar of Konrad-Adenauer-Stiftung



Simone Marcotullio
Italy

HIV/AIDS and Coinfections advocate for national and international communities



Marvelous Muchenje
Canada

Community Health Coordinator at Women's Health in Women's Hands Community Health Centre



Angelina Namiba
UK

Local and national strategy and policy adviser to ViiV, community representative; BHIVA Primary Care Working Group; S.A.F.E Kenya, Project Manager, Salamander Trust



Kneeshe Parkinson
USA

Patient Navigator/Co Chair Community Advisory Board (Part D) – Washington University School of Medicine in St. Louis/ Project ARK



Dr Bruno Spire
France

Senior Scientist at the French National Institute for Medical Research (INSERM)



Dr Andrew Ustianowski
UK

Consultant in Infectious Diseases and Tropical Medicine at the Pennine Acute Hospitals NHS Trust



Dr Benjamin Young
USA

SVP and CMO of the International Association of Providers in AIDS Care (IAPAC) and Editor of AIDSinfonet.org

Survey methodology*

The Positive Perspectives survey was undertaken in two phases:

1. A preliminary phase of qualitative research was carried out in June/July 2016 with 24 PLHIV, across four countries, to help understand the key areas of focus.
2. An in-depth quantitative survey was carried out in nine countries.


Participants were recruited in multiple ways, including: working with an expert panel (who supported recruitment via their own connections and networks), collaborating with charities, patient support organisations, non-government organisations and HIV online communities and promoting the research via social media:



PLHIV were eligible to enter the survey if they were over the age of 18 and diagnosed with HIV.

The screening process for participants involved completing a pre-questionnaire followed by a telephone interview.

The respondents below are broken down by country and sub-group.



*Further information on the methodology of the Positive Perspectives survey is available upon request.

Chapter 1: Diagnosis

Diagnosis is the first step of an important journey for PLHIV and is a step which many find challenging.⁵ Out of the 1,111 PLHIV surveyed, 83% (923) stated that their diagnosis had a negative impact on their emotional well-being.²

“I no longer have the motivation for things I used to do. I feel as though I have ‘lost myself’ and don’t know where to find it...”

Female
Aged 35–49
Canada

Chapter 1: Diagnosis





At the time of diagnosis, how do countries differ when looking at access to emotional support or guidance for PLHIV from their main doctor?^{1,2}



68%


(759)

of those surveyed indicated that they were offered support, referred for counselling or informed about support services by their main doctor **at diagnosis**



The results also show that those diagnosed over 10 years ago may have been offered less support at diagnosis (61%, 337)² than those diagnosed within the last two years (72%, 86).² Even with this positive change, the results suggested that further improvements still need to be made, with nearly a quarter (23%, 27)² of PLHIV who were diagnosed in 2015 to 2016 not being offered support, referred for counselling or informed about support services by their main doctor at diagnosis.

The Positive Perspectives survey results also highlighted that following diagnosis nearly half (45%, 502) of PLHIV turned to a close friend for emotional support,^{1,2} underlining the importance of personal relationships for those who are recently diagnosed.



of PLHIV (477) surveyed stated that being diagnosed with HIV has helped them to form more satisfying relationships²





Chapter 1: Diagnosis

The Positive Perspectives survey showed that 90% (998) of PLHIV actively sought support at diagnosis,² highlighting the importance of PLHIV being able to access the appropriate services and support networks at this time.

There were differences seen by country in those deciding not to seek support at diagnosis ranging from 2% (2) in Canada to 17% (20) in Italy.²

“ Reading other people’s experiences helps me to believe that I am going to be OK ”

Male
Aged 18–34
UK

“ My friends and family stand behind me even after the diagnosis and support me ”

Female
Aged 18–34
Germany

Chapter 1: Diagnosis

When looking at where PLHIV seek emotional support immediately following diagnosis we can see that:



45% (502)

of PLHIV from the above countries stated that **they would turn to a close friend for emotional support**²



54% (27)

of PLHIV in Austria chose their main HIV doctor as their number one choice for emotional support²

HIV patient support organisations are the number one choice for emotional support:²

36% (98)

For women



37% (109)

For heterosexuals



Compared with women, **more men seek emotional support from a close friend at diagnosis**²

MSM are more likely than heterosexuals to turn to someone they know living with HIV for emotional support²



51% (422)



vs

28% (76)



Chapter 1: Diagnosis

The respondents in the Positive Perspectives survey showed that how they gain emotional support today seems to have changed.

Of the 342 PLHIV surveyed that had a partner, **98%** (335) said that their partner knew about their HIV status:²

- **36%** (124) have HIV-positive partners²
- **63%** (215) have HIV-negative partners²
- **1%** (3) preferred not to say the HIV status of their partner²

"I am happy.
Life is smiling
on me"

Female
Aged 50+
Spain

What type of support do PLHIV turn to their partners for?



74%
(248)

Emotional support



52%
(174)

Reminds them to take
their medication



36%
(121)

Helps them to organise
their HIV medication



46%
(153)

Reminds them about
their HIV appointments



44%
(146)

Helps them to decide what
they need to discuss with their
main HIV healthcare provider



21%
(72)

Goes with them to routine doctor/
nurse appointments related to
their partner's HIV but does not sit
in the consultation itself



33%
(112)

Goes with them to routine doctor/
nurse appointments related to
their partner's HIV and sits in the
consultation itself

Source of emotional support today



From the four countries above, PLHIV are
most likely to turn to their main HIV provider
(35%, 255) for emotional support today²



Whilst PLHIV in Italy (31%, 38), Australia
(39%, 25) and Austria (38%, 19) are
most likely to still turn to a **close friend**
as they did at diagnosis²




PLHIV in Germany stated
that they are most likely
to turn to their **partner** for
emotional support
(32%, 45)²

Chapter 1: Diagnosis

HIV diagnosis has an emotional impact on people's lives. In the Positive Perspectives survey, PLHIV were asked to rate their emotional status on a scale from -50 to +50 at diagnosis and today:²

Mean rating of participants within the survey:



Chapter 2: Stigma and disclosure

Even though there have been advances in HIV treatment since the 1980s, the stigma associated with HIV still remains.

Participants in the survey were asked how their lives had changed both positively and negatively over the last five years of living with HIV. The top two negative responses reported by those surveyed were:²

1. "HIV still carries a lot of stigma"
2. "I am unable to engage in a proper/stable relationship"

"There is still
stigma with being
HIV positive. I still have
to hide my diagnosis "

Male
Aged 50+
UK



Chapter 2: Stigma and disclosure

The Positive Perspectives survey showed there were regional differences in how open PLHIV were in disclosing their status.¹

21%

in Europe
are generally
open
(127)

33%

in Australia
are generally
open
(21)

42%

in N. America
are generally
open
(184)


" Stigma surrounding
the disease is similar
to that of the 1990s "

Male
Aged 35–49
USA

93% of PLHIV have told their doctor (GP) about their HIV status:^{(1,034)²}


**#1
Reason:**

Wanted their doctor
to be involved in
their HIV care as
they believe that
they need to know
about all aspects of
their health²



**#1
Reason:**

Their doctor should
be aware of the HIV
medication they
are taking in case it
caused problems with
other medication they
are on²



PLHIV are not open about disclosing their HIV status to current work colleagues:



68%
(759)

of those surveyed have not disclosed
their status to their colleagues²



76%
(92)

of PLHIV in Italy have not disclosed
their status to their colleagues²

Chapter 2: Stigma and disclosure

82% (909) of PLHIV surveyed have experienced a form of stigma related to their HIV in the last 12 months,^{1,2} although this varies based on country, with 71% (86) of PLHIV in Italy having experienced recent stigma, and 90% (99) in Canada.²

Self-stigma



- Feelings of self-blame
- Guilt
- Lack of worth
- Need for secrecy

Institutionalised stigma



- Barred from jobs
- Being denied certain educational opportunities or health services

Physical stigma



- Being shunned or abandoned
- Harassment
- Asked to use separate facilities from others

Social stigma



- Feelings of isolation from my local community or social circle


Verbal stigma



- Gossip
- Taunting
- Scolding
- Labelling

“ I feel dirty and stigmatised. I have a lot of fear and loneliness ”

Female
Aged 35–49
Spain



Chapter 2: Stigma and disclosure

The Positive Perspectives survey results showed that self-stigma continues to have a big impact on the lives of PLHIV, with over a quarter (27%, 297) stating feelings of self-blame, guilt and a need for secrecy.^{1,2} Self-stigmatisation is reported more commonly in the UK (35%, 56) and Canada (33%, 36), but less so in Italy (19%, 23) and Austria (20%, 10).²


In relation to social stigma, nearly one in five (19%, 207) stated that they have feelings of isolation from their local community.^{1,2} Similarly, nearly a third (32%, 355) of PLHIV noted that their HIV status is only known to people very close to them and, even when asked, would not tell others.²

Chapter 2: Stigma and disclosure

The Positive Perspectives survey shows that:



(66) of those aged
18-34 are open
about their HIV
status²



(151) of those **>50**
years of age are
open about their
HIV status²

When comparing how stigma can vary from country to country, it can be seen that:

- **Across all the countries surveyed**, social stigma is the second-most dominant form of stigma raised by PLHIV²
- Participants from **North America** indicate that social stigma is almost, if not as big, of an issue as self-stigma²
- Participants from **Germany, Italy and Australia** feel as though they are very rarely impacted by social stigma²
- Participants from **Spain** are more than twice as likely to be impacted by self-stigma than social stigma²



Chapter 2: Stigma and disclosure

How to reduce feelings of stigmatisation:



67% ⁽⁶⁶¹⁾

of PLHIV from the countries above believe that better education for the general public was the number one way to address the issue²



62% ⁽⁷⁵⁾

believe that better teaching in schools was the number one way to address the issue²

A quarter ⁽²⁸³⁾ of those surveyed felt that better doctor education will reduce feelings of stigmatisation,^{1,2} which was a strong priority for those in Germany:²

These 283 PLHIV believe that additional education is required for:²



48% ⁽⁶⁷⁾



65% ⁽¹⁸⁴⁾

GPs & primary care doctors



64% ⁽¹⁸¹⁾

Hospital nurses



60% ⁽¹⁷¹⁾

Dentists

"Be careful he is sick, he has AIDS, he is toxic"

Male
Aged 35–49
Italy



Greater responsibility and better education of the mass media were also frequently selected as a way to combat stigma (44%, ⁴⁹⁴),^{1,2} particularly in Europe (50%, ³⁰⁷), less so in North America (35%, ¹⁵³).²

Chapter 3: Getting on treatment

The time to initiation of treatment for PLHIV has reduced over the last 10 years, with 88% ⁽¹⁰²⁾ ^{2,3} of those newly diagnosed, being on treatment within six months of diagnosis compared with 40% ⁽²¹⁸⁾ of those diagnosed >10 years ago. This equates to 52% ⁽⁵⁷¹⁾ across all PLHIV surveyed who have taken ART within six months of their HIV diagnosis.³ A key reason for this improvement is due to evolving treatment guidelines recommending earlier initiation of treatment.⁶

Today's treatments are effective in managing HIV and the latest survey findings suggest that PLHIV are optimistic about ongoing progress in treatment development, with 89% ⁽⁹⁸⁷⁾ feeling that advances in HIV treatment will improve their quality of life.^{1,2}

"These days I have
greater faith in the
research "

Female
Aged 50+
Italy


" Learning that I could
keep from passing the virus
on if I stay at non-detectable
levels was a game changer in
how I felt when seeking medical
care. I have returned to work part
time after being out of work for
nearly a decade because of the
advancements in treatment "

Male
Aged 35–49
USA



Chapter 3: Treatment satisfaction

Of the 98% (1,085) who are currently on treatment,⁴ 56% (604) are very satisfied with their treatment. This is quite consistent across gender, age and time since diagnosis.^{2,3} However, there are differences when looking at the results from various countries:²







of PLHIV in the US **are currently very satisfied** with their HIV treatment, which is the highest when compared with other countries.



of PLHIV in Canada **are very satisfied**

In Europe, **lower treatment satisfaction levels** are seen in Spain and Austria, whilst **higher levels** are seen in the UK and Germany.²






"If I experience side effects, I can speak to my doctor who will be able to change the treatment"

Female
Aged 50+
UK

Chapter 3: Treatment satisfaction

Around 3 in 10 PLHIV agree that their HIV treatment gives them unpleasant side effects, which were seen as the predominant reason for 43% (355) of those surveyed to switch their treatment.^{3,4} Other reasons for switching include helping to improve QoL (37%, 310) and to reduce the number of pills they need to take (32%, 264).²

When looking at how PLHIV made the decision to switch treatment, 55% (460) reported that the decision was jointly made between them and their main HIV doctor:³



said that it was solely or primarily their doctor's decision




of PLHIV said that it was solely their decision

Chapter 4:

Conversations between PLHIV and their doctor

While the majority (71% ⁽⁷⁸⁹⁾) of PLHIV reported being very comfortable with raising issues of concern with their main doctor, this was lower in newly-diagnosed individuals, with only 61% ⁽⁷³⁾ feeling very comfortable.²

Interestingly, when this is broken down to a regional and country level there are some significant differences, with 81% ⁽³⁵³⁾ very comfortable in North America³ versus 62% ⁽³⁸¹⁾ in Europe.² Within Europe, the lowest levels are seen in Italy with only 56% ⁽⁶⁸⁾ very comfortable and in Spain with 50% ⁽⁶⁶⁾.²



of those surveyed visited their HIV doctor at least four times in the past 12 months.

"I hate when it comes to taking medication and when I have to go to doctors' appointments "


Male
Aged 35–49
Canada





Chapter 5: Long-term effects of treatment

Although, 72% (783) of PLHIV worry about the long-term effects of HIV treatments, more than a quarter (26%, 206) have not discussed this concern with their doctor.³



of PLHIV that are newly diagnosed worry about the long-term effects of treatment²

“ I have been on medication for HIV since 1996 and do sometimes worry what the long-term effect this is having on my body. Although I feel well and healthy at the moment I was diagnosed with osteoporosis about 8 years ago, and so sometimes I think what else could I develop as I get older? ”

Male
Aged 50+
UK

Chapter 5: Long-term effects of treatment

Long-term effects from treatment appear to be a large worry for those living with HIV. When asked on how to mitigate the impact of long-term ART, those surveyed answered:³



13%

(103)

speak to their doctor for
support and information



19%

(147)

read up and educate
themselves about HIV

9%

(74)

change or plan to
change their treatment





References

- 1 Murungi A, et al. Experience of living with HIV: Diagnosis & Disclosure – findings from the Positive Perspectives study. Presented at the IAS Conference on HIV Science (IAS 2017), 23-26 July 2017, Paris, France. Abstract: WEPED1423
- 2 Positive Perspectives Survey, 2017 DOF
- 3 Young B, et al. Patient Experience & Views on Antiretroviral Treatment – Findings from the Positive Perspectives Survey. Presented at the Infectious Disease Week (ID Week conference), 4–8 October 2017, San Diego, USA. Abstract number #1393.
- 4 Marcotullio S, et al. EU Patient Experience & Views on Antiretroviral Treatment – Findings from the Positive Perspectives Survey. Presented at the 16th European AIDS Conference (EACS), 25–27 October 2017, Milan, Italy. Abstract number #PE25/9.
- 5 Moskowitz JT, et al. Illness Appraisals and Depression in the First Year after HIV Diagnosis. *PLoS ONE*. 2013;8(10):e78904.
- 6 Trickey A, et al. Survival of HIV-positive patients starting antiretroviral therapy between 1996 and 2013: a collaborative analysis of cohort studies. *Lancet HIV*. 2017 [http://www.thelancet.com/journals/lanhiv/article/PIIS2352-3018\(17\)30066-8/fulltext?elsca1=tlpr](http://www.thelancet.com/journals/lanhiv/article/PIIS2352-3018(17)30066-8/fulltext?elsca1=tlpr)

Stay tuned for the attitudes
and perspectives of the
partners/significant others
of those that live with HIV
– **Coming in 2018!**

About ViiV Healthcare

ViiV Healthcare is a global specialist HIV company established in November 2009 by GlaxoSmithKline (LSE: GSK) and Pfizer (NYSE: PFE) dedicated to delivering advances in treatment and care for PLHIV and for people who are at risk of becoming infected with HIV. Shionogi joined in October 2012. The company's aim is to take a deeper and broader interest in HIV/AIDS than any company has done before and take a new approach to deliver effective and innovative medicines for HIV treatment and prevention, as well as support communities affected by HIV. For more information on the company, its management, portfolio, pipeline and commitment, please visit www.viivhealthcare.com.

About Positive Perspectives survey

The Positive Perspectives survey was sponsored by ViiV Healthcare and conducted by GfK UK Limited, an independent market research agency. The survey was conducted in nine countries (Austria, Australia, Canada, France, Germany, Italy, Spain, UK and USA) with a total of 1,111 PLHIV. The recruitment of PLHIV for this survey included working with charities, patient support groups and NGOs, HIV online communities and also promoting the research via social media (Facebook / Instagram / Twitter) in each country. The survey questionnaire was designed by ViiV Healthcare in collaboration with GfK UK Limited and through consultation with an independent Steering Committee of HIV professionals and advocates. The Positive Perspectives survey data are owned by ViiV Healthcare and will be presented at upcoming conferences, subject to abstract acceptance. ©2017 ViiV Healthcare group of companies or its licensor.

Hear more about the Positive Perspectives survey on Twitter and LinkedIn:

